

NPK Solutions

Veget-8 | 16 - 8 - 24 + 3MgO + TE

Eurosolids WS NPKs are top quality blends made from the purest raw materials, with limited caking sensitivity and with high level EDTA-chelated micro-nutrients for fertigation or foliar application. Eurosolids Veget-8 | 16 - 8 - 24 + 3MgO + TE is destined to be used during periods of intensive vegetative growth and protein build up in grain or fruit.

A healthy green plant is crucial to secure adequate photosynthesis and high yields.

- Veget-8 supports vegetative development and canopy expansion.
- Micronutrients 100% chelated and fully available
- Developed for fertigation in open field and greenhouses. Also suitable for foliar application



Product Characteristics

- Dissolves rapidly
- Low in Sodium and Chloride
- Easy to handle, dissolve and apply
- Production process certified according to ISO 9001:2015

Compatibility

Compatible with other fertilizers, except Calcium-containing fertilizers. Therefore, a separate fertilizer dissolving tank is required or the fertilizers must be applied at different times. The pH of the tank solution should be above 4.


Composition [%w/w]


Total Nitrogen	16%
Nitric Nitrogen	7%
Ammoniacal Nitrogen	4.5%
Ureic Nitrogen	4.5%
Phosphorus Pentoxide	8%
Potassium Oxide	24%
Magnesium Oxide	3%
Sulphur Trioxide	15%
Boron	0.010%
Copper	0.006%
Iron	0.056%
Manganese	0.028%
Molybdenum	0.004%
Zinc	0.008%

Packaging

- 25kg
- 500kg

Application





NPK Solutions

Veget-8 | 16 - 8 - 24 + 3MgO + TE

Dosage | Fertigation

Crop	Application date	Min kg/ha/season	Max kg/ha/season
Rose	• During entire fertigation program	200	350
Brocoli	• Starting flower formation and until 2 weeks before harvest	100	250
Tomato	• With first production phase and until the end of fertigation cycle	100	300
Pepper	• With first production phase and until the end of fertigation cycle	100	300

Dosage | Foliar application

Crop	Application date	Min kg/ha/appl	Max kg/ha/appl	Conc. % [w/v]
Fruit trees	3 - 5 applications: • As of fruit setting • 14 - 21 days interval	7	12	1.5 - 2.5
Vineyards	• As of end of flowering • Until 2 - 4 weeks before harvesting	7	12	1.5 - 2.5
Fruit vegetables	2 - 4 applications: • As of fruit setting • 10 - 14 days interval	4	8	1.0 - 1.5
Leaf vegetables	• During vegetative growth • Until 2 - 4 weeks before harvesting	3	6	0.5 - 1.0

In case of foliar feeding as part of a mix with crop protection products or other fertilizers, a compatibility test has to be done prior to preparing the spray-mix.

The mentioned dosages and application stages are given as a guideline. Exact dosages, concentration and application stage are subject to local conditions, use of other fertilizers and can only be given after an objective diagnosis.